

# Journal of New Century Innovations

VOLUME  
**29**  
ISSUE-2



Journal of new  
century innovations

Exact and natural sciences



ISSN (p): 2181-3671  
ISSN (e): 2181-368X



[newjournal.org](http://newjournal.org)

OLMOS PANJARADAGI DISKRET SHRYODINGER  
OPERATORINING SPEKTRI

*Xayitova Xilola G'afurovna,*  
*Buxoro davlat universiteti*  
*«Matematik analiz» kafedrasi o'qituvchisi*  
*Ikromova Sarvinoz Ismoil qizi*  
*Buxoro davlat universiteti*  
*Fizika-matematika fakulteti magistri*

**Annotatsiya.** Ushbu maqolada olmos panjaradagi sistemada aniqlangan diskret Shryodinger operatori qaralgan. Olmos panjaradagi sistemada aniqlangan diskret Shryodinger operatori tavsivlangan hamda bu operatorning impuls tasviri olingan va muhim spektri o'rganilgan. Ma'lumki, olmos panjaradagi ikki zarrachali Shryodinger operatorlarini o'rganish ochiq masala hisoblanadi. Mazkur maqolada o'rganilayotgan operator olmos panjaradagi ikki zarrachali Shryodinger operatorini o'rganishda muhim ahamiyat kasb etadi.

**Kalit so'zlar:** diskret Shryodinger operatori, kompakt operator, muhim spektr, xos qiymat, xos funktsiya.

SPECTRUM OF THE DISCRETE SCHROEDINGER  
OPERATOR ON A DIAMOND LATTICE

Khayitova Khilola Gafurovna,  
Department of Mathematical Analysis  
Bukhara State University  
Ikromova Sarvinoz Ismoil qizi  
Master Student, Faculty of Physics and Mathematics,  
Bukhara State University

**Annotation.** This paper considers the discrete Schrödinger operator defined in a system on a diamond lattice. The discrete Schrödinger operator defined in the diamond lattice system is described, and the pulse image of this operator is obtained and its essential spectrum is studied. It is known that the study of two-particle Schrödinger operators in a diamond lattice is an open problem. The operator studied in this article is of great importance in the study of the two-particle Schrödinger operator in the diamond lattice.

**Keywords:** Discrete Schrödinger operator, compact operator, critical spectrum, eigenvalue, eigenfunction.

$\mathbb{C}$ – bir o'lchamli kompleks sonlar fazosi bo'lsin. Ixtiyoriy  $n \in \mathbb{N}$  natural soni uchun  $L_2[a, b]^n$  orqali  $[a, b]^n$  da aniqlangan kvadrati bilan integrallanuvchi (umuman olganda kompleks qiymatli) funksiyalarning Hilbert fazosini belgilaymiz.  $\mathbb{T}^d$  bilan  $d$  o'lchamli tor, ya'ni

$$\mathbb{T}^d = \underbrace{\mathbb{T}^1 \times \mathbb{T}^1 \times \dots \times \mathbb{T}^1}_{dmarta}$$

ni belgilaymiz.  $d$  o'lchamli tor  $\mathbb{T}^d$  da aniqlangan, Haar ma'nosida o'lchovga ega va

$$\int_{\mathbb{T}^d} |f(x)|^p dx < \infty$$

shartni qanoatlantiruvchi barcha  $f: \mathbb{T}^d \rightarrow \mathbb{C}$  funksiyalarning chiziqli fazosini qaraymiz, bunda integralda o'lchov Haar ma'nosida olinadi va  $p$  tayinlangan musbat son. Elementlarni qo'shish va songa ko'paytirish odatdagi funksiyalarni qo'shish va songa ko'paytirish kabi kiritiladi. Hosil bo'lgan fazo  $L_p(\mathbb{T}^d)$  kabi belgilanadi. Demak,  $L_p(\mathbb{T}^d)$  fazoning elementlari  $\mathbb{T}^d$  da aniqlangan va har bir o'zgaruvchisi bo'yicha  $2\pi$  davrga ega bo'lgan funksiyalardir [1-6].

Ushbu maqolada  $L_2^{(2)}(\mathbb{T}^2)$  – Hilbert fazosida  $H = Q + H_0$  ko'rinishda aniqlangan Olmos panjaradagi diskret Shredinger operatori  $H = H_0 + Q$  ning muhim spektrini o'rganamiz.

Quyidagi to'plamni kiritamiz:

$$A = \{v(n): v(n) = n_1 v_1 + n_2 v_2 \quad n = (n_1; n_2), n \in \mathbb{Z}^2\},$$

bu yerda

$$v_1 = e_3 - e_1 = (-1; 0; 1), \quad v_2 = e_3 - e_1 = (0; -1; 1).$$

**Ta'rif.**  $A$  to'plamga ikki o'lchamli olmos panjara deyiladi.

Quyidagi to'plamni kiritamiz:

$$\Omega = A \cup (p + A), \quad p = \frac{1}{3}(v_2 - v_1) = \frac{1}{3}(-1; -1; 2).$$

$\ell_2(\Omega)$  - orqali  $\Omega$  da kvadrati bilan jamlanuvchi  $\hat{f}(n) = (\hat{f}_1(n), \hat{f}_2(n))$  funksiyalar juftligini belgilaymiz. Bu fazo Hilbert fazosi bo'lib, skalyar ko'paytma quyidagicha aniqlangan:

$$(\hat{f}, \hat{g}) = \sum_{v \in A} 3\hat{f}_1(n)\hat{g}_1(n) + \sum_{v \in (p+A)} 3\hat{f}_2(n)\hat{g}_2(n).$$

$L_2^{(2)}(\mathbb{T}^2) - \mathbb{T}^2$  da aniqlangan kvadrati bilan integrallanuvchi  $f(x) = (f_1(x), f_2(x))$  funksiyalar juftligining Hilbert fazosi bo'lsin. Bu yerda skalyar ko'paytma quyidagicha aniqlangan

$$(f, g) = (f_1, g_1) + (f_2, g_2)$$

Bunda

$$(f_i, g_i) = \int_{\mathbb{T}^2} f_i(x)\overline{g_i(x)} dx, \quad i = 1, 2.$$

Quyidagi

$$\mathcal{F}: \ell_2(\Omega) \rightarrow L_2^{(2)}(\mathbb{T}^2)$$

unitar operatorni kiritamiz:

$$\mathcal{F} = \begin{pmatrix} \mathcal{F} & 0 \\ 0 & \mathcal{F} \end{pmatrix},$$

$$(\mathcal{F}\hat{f})(x) = \frac{\sqrt{3}}{2\pi} \sum_{n \in \mathbb{Z}^2} e^{i(x,s)} \hat{f}(s).$$

Bu operator teskarisi  $\mathcal{F}^{-1}: L_2^{(2)}(\mathbb{T}^2) \rightarrow \ell_2(\Omega)$  quyidagicha aniqlanadi:

$$\mathcal{F}^{-1} = \begin{pmatrix} \mathcal{F}^{-1} & 0 \\ 0 & \mathcal{F}^{-1} \end{pmatrix},$$

$$(\mathcal{F}^{-1}f)(s) = \frac{\sqrt{3}}{2\pi} \int_{\mathbb{T}^2} e^{-i(s,x)} f(x) dx.$$

bu yerda

$$(s, x) = s_1 x_1 + s_2 x_2.$$

Olmos panjaradagi diskret Shredinger operatori  $\widehat{H}$  ushbu  $\ell_2(\Omega)$  fazoda chegaralangan o‘z-o‘ziga qo‘shma operator sifatida quyidagicha aniqlanadi:

$$\widehat{H} = -3(\Delta_2 + 1) + \widehat{Q}.$$

Bunda

$$(-3(\Delta_2 + 1)\widehat{f})(v) = ((V_1\widehat{f}_2)(n); (V_2\widehat{f}_1)(n))$$

Bu yerda

$$(V_1\widehat{f}_2)(n) = \widehat{f}_2(n) + \widehat{f}_2(n - e_1) + \widehat{f}_2(n - e_2)$$

$$(V_2\widehat{f}_1)(n) = \widehat{f}_1(n) + \widehat{f}_1(n - e_1) + \widehat{f}_1(n - e_2)$$

$$e_1, e_2 \in \Omega, n = (n_1; n_2), e_1 = (1; 0), e_2 = (0; 1).$$

$\widehat{Q}$ -  $\Omega$  da aniqlangan zarrachalarning o‘zaro ta’sir potentsiali bo‘lib, ular quyidagi formulalar bilan aniqlanadi.

$$(\widehat{Q}f)(n) = \begin{pmatrix} \widehat{Q}_1(n) & 0 \\ 0 & \widehat{Q}_2(n) \end{pmatrix} \begin{pmatrix} \widehat{f}_1(n) \\ \widehat{f}_2(n) \end{pmatrix} = \begin{pmatrix} \widehat{Q}_1(n)\widehat{f}_1(n) \\ \widehat{Q}_2(n)\widehat{f}_2(n) \end{pmatrix}$$

bunda

$$\sum_{n \in A_2} |\widehat{Q}_1(n)| < \infty, \quad \sum_{n \in (p+A_2)} |\widehat{Q}_2(n)| < \infty.$$

$\widehat{H}$  operatorni koordinata ko‘rinishidan impuls tasvirga o‘tish  $F$  almashtirishilari yordamida amalga oshiriladi

$$H = \mathcal{F}\widehat{H}\mathcal{F}^{-1} = \mathcal{F}(-3(\Delta_2 + 1))\mathcal{F}^{-1} + \mathcal{F}\widehat{Q}\mathcal{F}^{-1}.$$

$H$  operator olmos panjaradagi diskret Shredinger operatorining impuls tasviri bo‘lib, u quyidagicha aniqlanadi  $H = H_0 + Q$ ,  $H_0$  va  $Q$  matritsa operatorlari bo‘lib,  $L_2^{(2)}(\mathbb{T}^2)$  da quyidagicha aniqlanadi

$$(H_0f)(x) = \begin{pmatrix} 0 & E(x) \\ E(x) & 0 \end{pmatrix} \begin{pmatrix} f_1(x) \\ f_2(x) \end{pmatrix} = \begin{pmatrix} E(x)f_2(x) \\ E(x)f_1(x) \end{pmatrix},$$

$$(Qf)(x) = \begin{pmatrix} Q_1 & 0 \\ 0 & Q_2 \end{pmatrix} \begin{pmatrix} f_1(x) \\ f_2(x) \end{pmatrix} = \begin{pmatrix} (Q_1f_1)(x) \\ (Q_2f_2)(x) \end{pmatrix},$$

Bunda,  $E(x) - 2$  o‘zgaruvchili kompleks qiymatli funksiya

$E(x) = \frac{1}{3}(1 + e^{ix_1} + e^{ix_2})$ ,  $Q_i - L_2(\mathbb{T}^2)$  da aniqlangan integral operator

$$(Q_i f_i)(x) = \int_{\mathbb{T}^2} Q_i(x-t) f_i(t) dt. \quad i = 1, 2,$$

$Q_i(\cdot) - \mathbb{T}^2$  da aniqlangan haqiqiy qiymatli biror uzluksiz, juft funksiya.

Quyidagi teoremani qaraymiz.

**Teorema.**  $\sigma(H_0) = [-1; 1]$ .

**Isbot.** Bizga ma'lumki  $H_0$  matritsaviy operatorining spektri quyidagi formula bilan aniqlanadi

$$\sigma(H_0) = \bigcup_{x \in \mathbb{T}^2} \sigma(H_0(x)). \quad (2)$$

Bunda  $H_0(x)$  – har bir fiksilangan  $x \in \mathbb{T}^2$  da ikki o'lchamli matritsa, ya'ni

$$H_0(x) = \begin{pmatrix} 0 & E(x) \\ \overline{E(x)} & 0 \end{pmatrix},$$

$$E(x) = \frac{1}{3}(1 + e^{ix_1} + e^{ix_2}).$$

Shuning uchun  $H_0(x)$  ning spektri xos qiymatlaridan iborat bo'ladi, ya'ni har bir tayinlangan  $x \in \mathbb{T}^2$  larda  $\det|H_0 - \lambda I| = 0$  tenglamaning ildizlaridan iboratdir. Ushbu tenglamani tuzamiz:

$$(H_0 - \lambda I)(x) = \begin{pmatrix} 0 & E(x) \\ \overline{E(x)} & 0 \end{pmatrix} - \begin{pmatrix} \lambda & 0 \\ 0 & \lambda \end{pmatrix} = \begin{pmatrix} -\lambda & E(x) \\ \overline{E(x)} & -\lambda \end{pmatrix}.$$

$$\det|H_0 - \lambda I| = \begin{vmatrix} -\lambda & E(x) \\ \overline{E(x)} & -\lambda \end{vmatrix} = 0,$$

$$\lambda^2 - E(x) \cdot \overline{E(x)} = 0,$$

$$\lambda_{1,2} = \pm |E(x)|, \quad x \in \mathbb{T}^2.$$

Bu yerda

$$|E(x)|^2 = E(x)\overline{E(x)} = \frac{1}{3}(1 + e^{ix_1} + e^{ix_2})\frac{1}{3}(1 + e^{-ix_1} + e^{-ix_2})$$

$$= \frac{1}{9}(3 + e^{ix_1} + e^{-ix_1} + e^{ix_2} + e^{-ix_2} + e^{i(x_1-x_2)} + e^{-i(x_1-x_2)}) =$$

$$= \frac{1}{9}(3 + 2 \cos x_1 + 2 \cos x_2 + 2 \cos(x_1 - x_2)).$$

Shunday qilib,

$$\sigma(H_0(x)) = \{-|E(x)|; |E(x)|\}.$$

Demak, (2) ga ko'ra

$$\sigma(H_0) = \bigcup_{x \in \mathbb{T}^2} \sigma(H_0(x)) = \bigcup_{x \in \mathbb{T}^2} \{-|E(x)|; |E(x)|\}$$

$$= -Ran\{|E(x)|\} \cup Ran\{|E(x)|\}.$$

Endi

$$\max_{x \in \mathbb{T}^2} |E(x)| = \max_{x \in \mathbb{T}^2} \frac{1}{9}(3 + 2 \cos x_1 + 2 \cos x_2 + 2 \cos(x_1 - x_2)) = 1$$

va

$$\min_{x \in \mathbb{T}^2} |E(x)| = 0$$

ekanligidan, ushbu

$$-Ran\{|E(x)|\} = [-1; 0]$$

va

$$Ran\{|E(x)|\} = [0; 1]$$

tengliklarni hosil qilamiz. Shunday qilib,

$$\text{Demak } \sigma(H_0) = [-1; 1].$$

Teorema isbotlandi [7-11].

**Lemma.**  $Q: L_2^{(2)}(\mathbb{T}^2) \rightarrow L_2^{(2)}(\mathbb{T}^2)$  kompakt operator.



**Isbot.**  $Q: L_2^{(2)}(\mathbb{T}^2) \rightarrow L_2^{(2)}(\mathbb{T}^2)$  operatorni ko‘rinishi quydagicha edi:

$$(Qf)(x) = \begin{pmatrix} (Q_1f_1)(x) \\ (Q_2f_2)(x) \end{pmatrix}$$

Bunda,

$$(Q_i f_i)(x) = \int_{\mathbb{T}^2} Q_i(x-t)f_i(t)dt. \quad i = 1, 2.$$

$Q_i(\cdot)$  –ikkala o‘zgaruvchi bo‘yicha ham  $\mathbb{T}^2$  da aniqlangan biror uzluksiz funksiya. Biz  $Q$  operatorni kompakligini ko‘rsatishimiz uchun har bir  $i \in \{1, 2\}$  da  $Q_i : L_2(\mathbb{T}^2) \rightarrow L_2(\mathbb{T}^2)$  operatorni kompakt ekanligini ko‘rsatamiz. Yuqoridagi lemmadan ma‘lumki,

$$(Q_i f_i)(x) = \int_{\mathbb{T}^2} Q_i(x-t)f_i(t)dt$$

operator kompakt bo‘lishi uchun

$$\int_{\mathbb{T}^2} \int_{\mathbb{T}^2} |Q_i(x-t)|^2 dt dx < \infty$$

bo‘lishi zarur va yetarli edi. Shartga ko‘ra  $Q_i(\cdot) - \mathbb{T}^2$  da aniqlangan biror uzluksiz funksiya.

Bundan

$$\int_{\mathbb{T}^2} \int_{\mathbb{T}^2} |Q_i(x-t)|^2 dt dx$$

integral mavjud va chekli. Demak  $Q_i$  kompakt, ya‘ni  $Q$  kompakt operator.

Quyidagi teorema  $H$  operatorning muhim spektri va  $H_0$  operatorning spektri o‘rtasidagi bog‘lanishni ifodalaydi.

**Teorema.**  $\sigma_{ess}(H) = \sigma(H_0) = [-1; 1]$ .

**Isbot.** Muhim spektr turg‘unligi haqidagi Veyl teoremasiga ko‘ra,

$$H = Q + H_0$$

operatorning muhim spektri  $Q$  kompakt qo‘zg‘alishda o‘zgarmaydi va  $H_0$  operator spektri bilan ustma-ust tushadi.  $Q$  kompakt operator. Bundan esa  $\sigma_{ess}(H) = \sigma(H_0) = [-1; 1]$  ekanligi kelib chiqadi. Teorema isbotlandi.

### FOYDALANILGAN ADABIYOTLAR

1. [T Rasulov](#), [N Tosheva](#). New branches of the essential spectrum of a family of 3x3 operator matrices. - Journal of Global Research in Math. Archive, 2019
2. Rasulov T.H., Tosheva N.A. (2019). Analytic description of the essential spectrum of a family of 3x3 operator matrices. Nanosystems: Phys., Chem., Math., 10:5, pp. 511-519.
3. Расулов Т.Х. (2016). О ветвях существенного спектра решетчатой модели спин-бозона с не более чем двумя фотонами. ТМФ, 186:2, С. 293-310.
4. Rasulov T.H., Dilmurodov E.B. (2019). Threshold analysis for a family of 2x2 operator matrices. Nanosystems: Phys., Chem., Math., 6(10), 616-622.

5. M.I.Muminov, T.H.Rasulov, N.A.Tosheva. Analysis of the discrete spectrum of the family of  $3 \times 3$  operator matrices. Communications in Mathematical Analysis. 23:1 (2020), pp. 17-37.

6. Т.Х.Расулов, Н.А.Тошева. О числе и местонахождении собственных значений обобщенной модели Фридрихса. Тезисы 42-й Всероссийской молодежной школа-конференции Современные проблемы математики. Екатеринбург, 2011, Стр. 102–104.

7. Бобоева М.Н. Поля значений одной  $2 \times 2$  операторной матрицы с одномерными интегральными операторами. Вестник науки и образования. 17-2 (95) (2020), С 14-18.

8. T.Rasulov, N.Tosheva. Main property of regularized Fredholm determinant corresponding to a family of  $3 \times 3$  operator matrices. European science. 2.(51) 2020, pp. 11-14

9. А.Алаудинов, Ch.Jumayeva. Local innerderivations on three-dimensional lie algebras. Fan va jamiyat. 2022 й. 2-сон. 5-8 б.

10. А.Алаудинов, Ch.Jumayeva. Local inner derivations on four-dimensional lie algebras. Қорақалпоғистонда фан ва таълим. 2022 й. 1 -сон. 22-29 б.

11. Сайлиева Г.Р. (2021). Использование новых педагогических технологий в обучении «Аналитическая геометрия». Вестник науки и образования. 68-71.