


ACADEMICIA

ISSN (online) : 2249-7137




ACADEMICIA

An International
Multidisciplinary Research
Journal



Published by
South Asian Academic Research Journals
A Publication of CDL College of Education, Jagadhri
(Affiliated to Kurukshetra University, Kurukshetra, India)

285.	PRELIMINARY STUDY OF SHAKYA BUDDHISTS Rudra Nath Dahal	1791-1799	10.5958/2249-7137.2021.00291.3
286.	FORMATION OF CIVIL IDENTITY OF UZBEKISTAN YOUTH AS A FACTOR OF INTERNATIONAL CONSENT Shakirdzhan Mamazhanovich Zakirov	1800-1804	10.5958/2249-7137.2021.00285.8
287.	ROLE OF THE DESIGN CODE IN IMPROVING THE QUALITY OF THE URBAN ENVIRONMENT Shukhrat Reyimbaev, Zarif Adilov, Zafar Matniyozov	1805-1812	10.5958/2249-7137.2021.00286.X
288.	PEDAGOGICAL ISSUES IN ENSURING CONTINUOUS AND INTERMITTENT STATE OF THE FIELD OF FINE ARTS IN THE SYSTEM OF ART EDUCATION U. Nurtaev	1813-1821	10.5958/2249-7137.2021.00287.1
289.	METHODS OF WORD FORMATION IN UZBEK DIALECTS OF THE REPUBLIC OF KARAKALPAKSTAN Umida Ibragimova	1822-1828	10.5958/2249-7137.2021.00288.3
290.	PHYSICS - MECHANICAL PROPERTIES OF IRRIGATED MEADOW SOILS IN BUKHARA REGION Nazarova Sevara Mustakimovna, Turayeva Nargiza Nematilloevna, Tursunov Ulugbek Yusuf ugli, Mamurova Maftuna Otabek kizi	1829-1834	10.5958/2249-7137.2021.00289.5
291.	HEALTHY LIFESTYLE Anvar Narzulloevich Asadulloev, Nurbibi Rakhimovna Ochilova, Oygul Gaparovna Jabbarova	1835-42	10.5958/2249-7137.2021.00290.1
292.	THE USE OF INFORMATION TECHNOLOGY AS MEANS OF MOTIVATING STUDENTS TO LEARNING A FOREIGN LANGUAGE Daminova N.R, Daminova M.Z	1843-1847	10.5958/2249-7137.2021.00294.9



DOI: **10.5958/2249-7137.2021.00289.5**

PHYSICS - MECHANICAL PROPERTIES OF IRRIGATED MEADOW SOILS IN BUKHARA REGION

Nazarova Sevara Mustakimovna*; **Turayeva Nargiza Nematilloevna****;
Tursunov Ulugbek Yusuf ugli***; **Mamurova Maftuna Otabek kizi*****

*Doctor of Philosophy (PhD),
Agricultural Sciences Bukhara State University,
UZBEKISTAN

**Senior Lecturer of Bukhara State University,
UZBEKISTAN

***Students of Bukhara State University,
UZBEKISTAN

ABSTRACT

The article provides data on the physical and mechanical properties of irrigated meadow soils of the Jandar region. Depending on the mechanical composition, the degree of salinity and cultivation, it varies within a wide range of physical and mechanical properties; certain differences are found caused by their genesis and regional characteristics.

KEYWORDS: *Mechanical Composition, Hardness, Density, Specific Gravity, Soil Porosity, Humus, Solid Residue, Nitrogen, Phosphorus, Potassium.*

INTRODUCTION

A number of scientific studies are being carried out in the Republic aimed at further developing agriculture, preserving, reproducing, increasing soil fertility, efficiently using land resources, optimizing the ecological state, assessing the water-physical, technological, agrochemical properties and the reclamation state of soils.

In this plan, special attention is paid to the development of agrotechnical, agrophysical measures, taking into account soil and climatic conditions, to the widespread use of scientific and practical achievements in improving, restoring and increasing soil fertility. "Population growth over the next 35 years will require an increase in food production of about 60%." Development of scientifically based measures aimed at increasing soil fertility. Improving the efficiency of

irrigated lands, protecting the soil cover from degradation processes and preventing them is one of the urgent tasks.

This study was carried out in accordance with the priority direction of the development of science and technology in the republic. Irrigated soils, common in the Jandar region, differ in their properties and characteristics not only in soil zones, but also in soil and climatic conditions. On the problem of studying soil fertility, its management and other properties, large-scale scientific research has been carried out both abroad and in our republic. Irrigated soils of the Bukhara oasis not only differ in the properties of the steppe zone, they differ in climatic districts.

The morphogenetic structure, geographical distribution, reclamation state, agrophysical and agrochemical properties of the soils of the Bukhara oasis and other regions have been studied by many scientists, such as K. Gafurov, S. A. Abdullaev [3.1982; 130 p], U. Tojiev [9.2004; 158-159 p.], X.T.Artikova [1.2005; 28 p. 2.2019; 62 p.], M.A. Mazirov, S.V. Makarychev [6.2018; 605 p.], R. Kurv antaev [4. 2000; 40 p. 5.2019; 91-95 p.], Nazarova S.M. [7. 2016; 60-66 p. 8.2018; 187-190 P.], Sharipov.O.B., Gafurorova L.A. [10.2018; 76-79-pp.], Hakimova N., Kurvantaev R. [11. 2020; 68-71 pp.] and others. However, scientific research on the study of the current meliorative state, physical and mechanical properties of irrigated meadow soils in the Jandar region has not been carried out sufficiently.

The aim of the research is to develop recommendations for the correct organization of soil cultivation by determining the physical and mechanical properties and assessing the reclamation state of irrigated meadow soils common in the Jandar region.

The studies were carried out in soil-field and analytical-laboratory conditions. The reliability of the data obtained was carried out using the Microsoft Excel program on the basis of the "Methodology of field experience" by B.A. Dospekhov.

Research results

The irrigated meadow soils of the Jandar region are heavy and medium loamy in terms of the content of water-soluble salts, not saline (dense residue 0.150-0.375%), in some places (mainly chlorine) slightly saline (0.014-0.031%).

In the studied soils, the humus content in the arable and subsoil layers is 0.94-0.63%. At the same time, in the lower layers, no sharp differences are observed in the humus content, and along the sections, the humus content is 0.41–0.30%.

The influence of the age of irrigation on the content of nutrient reserves (nitrogen, phosphorus and potassium) is clearly seen. In the irrigated meadow soils of the Jandar region, nitrogen is 1.9–3.2 t / ha, phosphorus is 6.5–14.5 t / ha, and potassium is 51.4–106.5 t / ha.

Physic mechanical properties of irrigated soils in the Jandar District It is noted that the soils are distinguished by their mechanical composition by their originality in the administrative and geomorphologic regions formed on alluvial deposits of the lower part of the river. Zarafshan [5. 2019; 91-95 p. 7.2016; 60-66 p. 8. 2018; 187-190 s.]. Basically, the mechanical composition consists of the following particles: coarse sand (1–0.25 mm), medium sand (0.25–0.1 mm), fine sand (0.1–0.05 mm).

The mechanical composition of the soils of the Jandar region consists of sandy, sandy loam, light, medium, heavy loams. The figure shows a map of the mechanical composition of the soils

of the Jandar region of the Istiklol (Scale 1: 10000), sandy loam, light-, medium-, heavy loamy (figure).

The specific gravity is a stable unit, it depends on the chemical, mineralogical composition and on the supply of humus to the soil. In irrigated meadow soils in the lower reaches of the Zarafshan in the district, the specific weight is 2.58–2.66 g / sm³.

Volumetric mass is a variable and different unit depending on.

Picture. Map of the mechanical composition of soils in the Jandar region of the Istiklol.

Various processes occurring in the soil. In the top, arable layer soil, the bulk density varies between 1.27–1.63 g / sm³ depending on humus content, texture, salinity and other properties. Among the upper layers, the highest density (1.53–1.63 g / sm³) is observed in the soils of the Jandar region.

Old-irrigated meadow soils, as a result of numerous visits of heavy machinery and non-compliance with the irrigation regime, increases the optimal density of the addition [4. 2000; 40-p.]. It was determined that under the conditions of irrigated meadow soils, their density increases with the weighting of the mechanical composition (Table 1).

In irrigated soils, the movement of water, the content of soluble salts, the preservation of moisture, and the provision of air to the root system is directly related to the porosity of the soil. In the studied meadow soils, the total porosity, depending on the duration of irrigation, varies widely across the genetic layers of the profile (42–51%).

TABLE 1 GENERAL PHYSICAL PROPERTIES OF IRRIGATED MEADOW SOILS OF THE JANDAR REGION

Cutting depth, sm	0-35	35-50	50-75	75-105	105-132	132-180
Specific weight, g / sm ³	2,58	2,67	2,64	2,60	2,66	2,66
Bulk weight, g / sm ³	1,34	1,49	1,53	1,52	1,53	1,48
Total porosity, %	48	44	43	42	42	45

Soil hardness is a technological indicator of a property, it is of particular importance in fertility and soil. The hardness of the genetic soil layers in the Jandar region, depending on the mechanical composition, layer moisture and density, varies from 5.1 to 16.7 kg / sm², high hardness indicators are confirmed by the data obtained for the subsoil layers (Table 2). The resisting density for the development of plant roots in deep layers is determined by the penetrometer device, the results obtained show in the Jandar region along the vertical horizons is 72.5–350 kPa.

The greatest resistance is observed in the layer 105-132 sm (350 kPa). The soils widespread in various farms resist the development of the root system to varying degrees, especially there is a high resistance for plant roots of common soils in the Istiklal.

TABLE 2 INDICATORS OF HARDNESS OF MEADOW SOILS OF JANDAR DISTRICT

Horizon depth, sm	Horizontal hardness, kg / sm ²	Vertical depth, sm	Vertical hardness, kPa
0-35	7,8	10	72,5
35-50	16,7	20	155
50-75	6,1	30	232
75-105	5,8	40	237
105-132	5,1	50	350
132-180	5,7	60	300
		70	250

The content of agronomic valuable macro-aggregates in the arable layer of old-irrigated medium loamy meadow soils in the Jandar region in the Istiklal is 52-73%.

V. CONCLUSION

Distributed in the lower part of the Bukhara oasis of soils, peculiar basic morphological features were formed in the process of prolonged irrigation. Soils of geomorphological regions are characterized by their peculiarity in the mechanical composition of soils and consist of sandy loam, light-, medium-, heavy-loamy varieties, mainly consisting of the following fractions: coarse, medium fine sand

The specific gravity in the genetic layers varies in the range of 2.56–2.67 g / sm³. Soils on the genetic horizons have different density (1.27-1.63 g / sm³), the total porosity in the upper layer is satisfactory (47-51%), in the lower layers it belongs to the unsatisfactory (38-42%) categories.

For genetic horizons, depending on the density and texture, the value of hardness varies from 5.1 to 16.7 kg / sm², soil support for root development is 72.5 - 350 kPa.

REFERENCES:

1. Artikova Kh.T. Soils of the Bukhara oasis and their change under the influence of irrigation. Abstract. Cand. diss. Tashkent. 2005 . 28 p.
2. Artikova X.T. Evolution, ecological state and soil fertility of the Bukhara oasis. Avoref. diss. Doctor of Science (DSc). Tashkent, 2019.-62 p.
3. Gafurov K., Abdullaev S. Characteristics of the soil cover of the irrigated zone of the Bukhara region. Tashkent: Publishing house "Fan", 1982. - 130 p.
4. Kurvantaev R. Optimization and regulation of the agrophysical state of irrigated soils in the desert zone of Uzbekistan. Author. doct. diss. Tashkent. 2000 .40p.
5. Kurvantaev R., Nazarova S.M. Agrophysical characteristics of irrigated meadow soils of the Bukhara oasis. // Modern trends in scientific support of the agro-industrial complex: Collective monograph. [editorial board: L.I.Ilyin and others; otv for issue VV Ogorkov]. Ivanovo, 2019 .91-95pp.

6. Mazirov M.A. S.V. Makarychev Chemical properties of irrigated and rainfed gray soils of the western Tien Shan. // Modern trends in scientific support of the agro-industrial complex of the Upper Volga region. Collective monograph. Moscow. Publishing house "Ivanovo", 2018. - 605 p.
7. Nazarova S.M., Kungirov H.N., Kurvantaev R.K., Dadamukhamedova M.R. The main factors of soil formation and their evolution in the Zerafshan valley. // In the collection: Ecological condition of the natural environment and scientific and practical aspects of modern reclamation technologies Collection of scientific papers. Ryazan, 2016.60-66pp.
8. Nazarova SM, Kurvantaev R. Mechanical composition of irrigated soils of the Bukhara oasis. / Actual problems of modern science. Scientific journal No. 4 (101), 2018. 187-190pp.
9. Tozhiev U.T., Nafetdinov Sh.Sh. Systemic research of diagnostics, classification and structure of soil cover in the Republic of Uzbekistan. 2004.153-159pp.
10. Sharipov.O.B., Gafurorova L.A. Biological activity of irrigated grassland alluvial soils of the Bukhara Oasis. / Evropean Science Review. Avstria, 2018. 76-79pp.
11. Hakimova N., Kurvantaev R. Evolution of raflux soils of the midrange of the valley Zerafshan / ANNALI D / LTALIA Scientific Journal of Italy, VOL. 2. 2020 (4). 68-71pp.
12. Атоева М.Ф. Периодичность обучения физике. Аспирант и соискатель. – Москва, 2010. – №6. – С. 41-43.
13. M.F. Atoyeva. Interdisciplinary relations in physics course at specialized secondary education. The Way of Science. – Volgograd, 2016. – №9 (31). – P.22-24.
14. M.F. Atoyeva. The significance of periodicity at teaching physics. The Way of Science. – Volgograd, 2016. – № 10 (32). – P.62-64.
15. Атоева М.Ф. Эффективность обучения электродинамике на основе технологии периодичности. The Way of Science. – Volgograd, 2016. – № 10 (32). – P.65-66.
16. M.F. Atoyeva. Use of Periodicity in Teaching Physics. Eastern European Scientific Journal. – Düsseldorf-Germany, 2017. № 4. –P. 35-39.
17. M.F. Atoyeva. Didactic foundations of inter-media relations in the training of university students. International Scientific Journal. **Theoretical & Applied Science**. p-ISSN: 2308-4944 (print) e-ISSN: 2409-0085 (online). **Year: 2020 Issue: 06 Volume: 86**, P. 124.
18. M.F. Atoyeva, R. Safarova. Pedagogical integration as a means of forming professionally important qualities among students of a medical university. *Academicia*. ISSN: 2249-7137 Vol. 10, Issue 8, August 2020. Impact Factor: SJIF 2020 = 7.13 *ACADEMICIA: An International Multidisciplinary Research Journal* <https://saarj.com>.
19. M.F. Atoyeva. Pedagogical Tests As An Element Of Types Of Pedagogical Technologies. *The American Journal of Applied Sciences*, 2(09), (TAJAS) SJIF-5.276 DOI-10.37547/tajas Volume 2 Issue 9, 19.09.2020. ISSN 2689-09. 92 **The USA Journals, USA** www.usajournalshub.com/index.php/tajas 164-169. Имп.**5.2**.

-
- 20.** Farkhodovna, A. M. (2020). The problems of preparing students for the use of school physical experiment in the context of specialized education at secondary schools. *European Journal of Research and Reflection in Educational Sciences*, 8 (9), 164-167.
- 21.** Saidov S.O., Fayzieva Kh. A., Yuldosheva N. B. Atoyeva M.F. The Elements Of Organization Of The Educational Process On The Basis Of New Pedagogical Technologies. *The American Journal of Applied Sciences*, 2(09), (TAJAS) SJIF-5.276 DOI-10.37547/tajas Volume 2 Issue 9, 19.09.2020. ISSN 2689-09.92 **The USA Journals, USA** www.usajournalshub.com/index.php/tajas 164- 169. **Имп.5.2.**
- 22.** Atoeva Mehriniso Farhodovna, Arabov Jasur Olimboevich, Kobilov Bakhtiyor Badriddinovich. (2020). Innovative Pedogogical Technologies For Training The Course Of Physics. *The American Journal of Interdisciplinary Innovations and Research*, 2(12), 82-91.
- 23.** Камолиддин Аъзамович Самиев, Курбон Сайфуллоевич Саидов, Алижон Аминов. Теоретическое исследование процессов тепло-и массообмена в солнечных опреснительных установках. Молодой ученый Общество с ограниченной ответственностью Издательство.Москва. 2015 год. №10. Стр 300-305.