

Научный вестник Бухарского государственного университета
Scientific reports of Bukhara State University

1/2023



@buxdu_uz



@buxdu1



@buxdu1



www.buxdu.uz



BUXORO DAVLAT UNIVERSITETI ILMIY AXBOROTI



1/2023

BUXORO DAVLAT UNIVERSITETI ILMIY AXBOROTI
SCIENTIFIC REPORTS OF BUKHARA STATE UNIVERSITY
НАУЧНЫЙ ВЕСТНИК БУХАРСКОГО ГОСУДАРСТВЕННОГО УНИВЕРСИТЕТА

Ilmiy-nazariy jurnal
2023, № 1

Jurnal 2003-yildan boshlab **filologiya** fanlari bo'yicha, 2015-yildan boshlab **fizika-matematika** fanlari bo'yicha, 2018-yildan boshlab **siyosiy** fanlar bo'yicha O'zbekiston Respublikasi Vazirlar Mahkamasi huzuridagi Oliy attestatsiya komissiyasining dissertatsiya ishlari natijalari yuzasidan ilmiy maqolalar chop etilishi lozim bo'lgan zaruriy nashrlar ro'yxatiga kiritilgan.

Jurnal 2000-yilda tashkil etilgan.
Jurnal 1 yilda 6 marta chiqadi.

Jurnal O'zbekiston matbuot va axborot agentligi Buxoro viloyat matbuot va axborot boshqarmasi tomonidan 2020-yil 24-avgust № 1103-sonli guvohnoma bilan ro'yxatga olingan.

Muassis: Buxoro davlat universiteti

Tahririyat manzili: 200117, O'zbekiston Respublikasi, Buxoro shahri Muhammad Iqbol ko'chasi, 11-uy.
Elektron manzil: nashriyot_buxdu@buxdu.uz

TAHRIR HAY'ATI:

Bosh muharrir: Xamidov Obidjon Xafizovich, iqtisodiyot fanlari doktori, professor

Bosh muharrir o'rinbosari: Rasulov To'liqin Husenovich, fizika-matematika fanlari doktori (DSc), professor

Kuzmichev Nikolay Dmitriyevich, fizika-matematika fanlari doktori (DSc), professor (N.P. Ogaryov nomidagi Mordova milliy tadqiqot davlat universiteti, Rossiya)

Danova M., filologiya fanlari doktori, professor (Bolgariya)

Margianti S.E., iqtisodiyot fanlari doktori, professor (Indoneziya)

Minin V.V., kimyo fanlari doktori (Rossiya)

Tashqarayev R.A., texnika fanlari doktori (Qozog'iston)

Mo'minov M.E., fizika-matematika fanlari nomzodi (Malayziya)

Adizov Baxtiyor Rahmonovich, pedagogika fanlari doktori, professor

Abuzalova Mexriniso Kadirovna, filologiya fanlari doktori, professor

Amonov Muxtor Raxmatovich, texnika fanlari doktori, professor

Barotov Sharif Ramazonovich, psixologiya fanlari doktori, professor, xalqaro psixologiya fanlari akademiyasining haqiqiy a'zosi (akademigi)

Baqoyeva Muhabbat Qayumovna, filologiya fanlari doktori, professor

Bo'riyev Sulaymon Bo'riyevich, biologiya fanlari doktori, professor

Djurayev Davron Raxmonovich, fizika-matematika fanlari doktori, professor

Durdiyev Durdimurod Qalandarovich, fizika-matematika fanlari doktori, professor

Olimov Shirinboy Sharofovich, pedagogika fanlari doktori, professor

Qahhorov Siddiq Qahhorovich, pedagogika fanlari doktori, professor

Umarov Baqo Bafoevich, kimyo fanlari doktori, professor

Murodov G'ayrat Nekovich, filologiya fanlari doktori, professor

O'rayeva Darmonoy Saidjonovna, filologiya fanlari doktori, professor

Navro'z-zoda Baxtiyor Nigmatovich, iqtisodiyot fanlari doktori, professor

Hayitov Shodmon Ahmadovich, tarix fanlari doktori, professor

To'rayev Halim Hojiyevich, tarix fanlari doktori, professor

Rasulov Baxtiyor Mamajonovich, tarix fanlari doktori, professor (Andijon davlat Pedagogika instituti rektori)

Boboyev Feruz Sayfullayevich, tarix fanlari doktori (O'ZR FA tarix instituti yetakchi ilmiy xodimi)

Jo'rayev Narzulla Qosimovich, siyosiy fanlar doktori, professor

Qurbonova Gulnoz Negmatovna, pedagogika fanlari doktori (DSc), professor

Jumayev Rustam G'aniyevich, siyosiy fanlar nomzodi, dotsent

Quvvatova Dilrabo Habibovna, filologiya fanlari doktori, professor

Axmedova Shoir Nematovna, filologiya fanlari doktori, professor

Amonova Zilola Qodirovna, filologiya fanlari doktori (DSc), dotsent

Zaripov Gulmurot Toxirovich, texnika fanlari nomzodi, dotsent

MUNDARIJA *** СОДЕРЖАНИЕ *** CONTENTS		
ANIQ VA TABIIY FANLAR *** EXACT AND NATURAL SCIENCES *** ТОЧНЫЕ И ЕСТЕСТВЕННЫЕ НАУКИ		
Rasulov T.H., Dilmurodov E.B.,	Kompakt qo'zg'alishli umumlashgan Fridriks modelining ba'zi spektral xossalari	4
Ismoilova D.E.	Fermionli fok fazodagi uchinchi tartibli operatorli matritsaga mos kanal operator va uning spektri	9
Jumaev J.J., Tursunova S.F., Ibragimova Sh.E.,	Inverse coefficient problem for heat equation in the bounded domain	15
Djurayev D.R., Turayev A.A., To'rayev O.G'.	Bi _{1.7} Pb _{0.3} Sr ₂ Ca _n Cu _{n-1} O _y vismut asosli kupratlarning olinish texnologiyasi	21
Umarov Sh.A.	Shifrlash algoritmlarida kriptobardoshli mantiqiy amallardan foydalanish usullari	25
Axmedov O.C.	Свободные и вынужденные осесимметричные колебания систем вязкоупругих цилиндрических оболочек	30
Исломов Б.И., Жураев Ф.М.	Краевые задачи с условием Геллерстедта на параллельных характеристиках для вырождающегося нагруженного уравнения параболо-гиперболического типа	34
Амиров С.Ф, Шаропов Ш.А, Сатторов Т.А.	Дифференциал трансформатор датчиклар оддий ва махсус структурали нозичиқ магнит занжирларининг математик моделлари	43
Турдиев Х.Х., Суяров Т.Р., Наврүзова М.Н.	Начально-краевая задача для системы интегро-дифференциальных гиперболических уравнений	50
Tosheva N.A.	Lower bound of the essential spectrum of a family of 3×3 operator matrices	57
Akramova D.I.	Estimates for convolution operators related to A_∞ type singularities	63
Jo'raqulova F.M.	Bozonli fok fazodagi operatorli matritsaga mos Faddeyev tenglamasi	72
Rasulov T.H., Ne'matova Sh.B.	Umumlashgan Fridriks modeli kvadratik sonli tasvirining komponentlari	77
Seytov Sh.J., Ochilova G.B.	Баъзи даврий реакцияларнинг математик моделлари	82
Sirliyeva F.A., Khudaybergenov K.K., Muminov Z.E.	Fuzzy neural network and agent technologies in the structural-parametric modeling of technological systems	87
Tosheva N. A., Qodirov S. O.	Python dasturlash tili yordamida matritsa sonli tasvirini kompleks tekislikda tasvirlash algoritmi	93
Хасанов И.И., Хасанов К.Х.	Начально-краевая задача для адвекции - дисперсионного уравнения дробного порядка	100
Husenov B. E.	Poisson representation for $A(z)$ -harmonic functions belonging to some class	107
TILSHUNOSLIK *** LINGUISTICS *** ЯЗЫКОЗНАНИЕ		
Nigmatova L. X.	Lisoniy sistemaning pog'onaviyligi va unda gipo-giperonimik munosabatlar	116
Sadigova S.I.	Phraseological combinations with ordinal numbers in English	123
Abdulxayrov D.P.	-Gan ekan shaklining mazmuniy ko'rinishlari	127
Xodjayeva D.I., Hoshimova G.M.	Erkin Vohidov she'rlarida ritorik so'roq gaplarning o'ziga xos xususiyatlari	131

LOWER BOUND OF THE ESSENTIAL SPECTRUM OF A FAMILY
OF 3×3 OPERATOR MATRICES**Tosheva Nargiza Ahmedovna,**

Bukhara State University, Bukhara, Uzbekistan

n.a.tosheva@buxdu.uz

Abstract. A family of 3×3 operator matrices $H(K)$, $K \in (-\pi, \pi]$ corresponding to the system of non-conserved and no more than three particles on a 1D lattice is considered. Considered family of operator matrices is defined as a linear, bounded and self-adjoint operator acting in the three-particle cut subspace of the Fock space. The number and location of the eigenvalues of the generalized Friedrichs model is investigated. We find a finite set $\Lambda \subset (-\pi, \pi]$ and give an estimate for the lower bound of the essential spectrum of operator matrix $H(K)$ for all $K \in \Lambda$.

Keywords: operator matrix; bosonic Fock space; annihilation and creation operators; generalized Friedrichs model.

Аннотация. Рассматривается семейство 3×3 операторных матриц $H(K)$, $K \in (-\pi, \pi]$, соответствующих системам с несохраняющимся и не более трёх частиц на одномерной решётке. Исследуемое семейство операторных матриц определено как линейный, ограниченный и самосопряжённый оператор в трёхчастичном обрезанном подпространстве фоковского пространства. Исследовано число и местонахождение собственных значений обобщённой модели Фридрихса. Найдено конечное множество $\Lambda \subset (-\pi, \pi]$, и получена оценка для нижней грани существенного спектра операторной матрицы $H(K)$ при всех $K \in \Lambda$.

Ключевые слова: операторная матрица, бозонное пространство Фока, операторы уничтожения и рождения, обобщённая модель Фридрихса.

Annotatsiya. Bir o'lchamli panjaradagi soni saqlanmaydigan va uchmadan oshmaydigan zarrachalar sistemasiga mos $H(K)$, $K \in (-\pi, \pi]$ – 3×3 operatorli matritsalar oilasi qaralgan. O'rganilayotgan operatorli matritsalar oilasi Fok fazosining qirg'ilgan uch zarrachali qism fazosidagi chiziqli, chegaralangan va o'z-o'ziga qo'shma operator sifatida aniqlangan. Umumlashgan Fridriks modeli xos qiymatlari soni va joylashuv o'rni tadqiq qilingan. Shunday $\Lambda \subset (-\pi, \pi]$ chekli to'plam topilib, barcha $K \in \Lambda$ larda $H(K)$ operatorli matritsa muhim spektrining quyi chegarasi uchun baholash olingan.

Kalit so'zlar: operatorli matritsa; bozonli Fok fazosi; yo'qotish va paydo qilish operatorlari; umumlashgan Fridriks modeli

Introduction. Block operator matrices are matrices the entries of which are linear operators between Banach or Hilbert spaces [1, 2]. Every bounded linear operator can be written as a block operator matrix if the space in which it acts is decomposed in two or more more components. Operator matrices arise in various areas of mathematics and its applications: in system theory as Hamiltonians, in the discretization of partial differential equations as large partitioned matrices due to sparsity patterns, in saddle point problems in non-linear analysis, in evolution problems as linearizations of second order Cauchy problems and as linear operators describing coupled systems of partial differential equations. Such systems occur widely in mathematical physics, e.g. in fluid mechanics, magnetohydrodynamics, theory of solid-state physics, quantum field theory, statistical physics and quantum mechanics.

Notice that one special class of block operator matrices are Hamiltonians associated with systems of non-conserved number of quasi-particles on a lattice. Their number can be unbounded as in the case of spin-boson models or bounded as in the case of "truncated" spin-boson models. It is remarkable that lattice spin-boson model with at most two photons can be represented as a 6×6 operator matrix which is unitarily equivalent to a 2×2 block diagonal operator with the 3×3 operator matrices on the diagonal [3,4].

In the present paper we consider a family of 3×3 operator matrices $H(K)$, $K \in T := (-\pi; \pi]$. These operator matrices are associated with the lattice systems describing two identical bosons and one particle, another nature in interactions, without conservation of the number of particles. They act in the direct

sum of zero-, one- and two-particle subspaces of the bosonic Fock space. We discuss the case where the dispersion function $\varepsilon(\cdot)$ has the form $\varepsilon(x) = 1 - \cos(nx)$ with $n > 1$. We denote by $\mathbb{T}$ the set of points T where the function $\varepsilon(\cdot)$ takes its (global) minimum. Under some smoothness assumptions on the parameters of $H(K)$, we estimate the lower bound of the essential spectrum of $H(K)$.

Family of 3×3 operator matrices and main results

Let $\mathbb{T}$ be the one-dimensional torus, $H_0 := \mathbb{C}$ be the field of complex numbers, $H_1 := L_2(\mathbb{T})$ be the Hilbert space of square integrable (complex) functions defined on $\mathbb{T}$ and $H_2 := L_2^s(\mathbb{T}^2)$ be the Hilbert space of square integrable (complex) symmetric functions defined on $\mathbb{T}^2$. The Hilbert space $H := H_0 \oplus H_1 \oplus H_2$ is called three-particle cut subspace of a bosonic Fock space $F_s(L_2(\mathbb{T}))$ over $L_2(\mathbb{T})$, respectively.

In the present paper we consider a family of 3×3 operator matrices $H(K)$, $K \in \mathbb{T}$ acting in the Hilbert space H as

$$H(K) := \begin{pmatrix} H_{00}(K) & H_{01} & 0 \\ H_{01}^* & H_{11}(K) & H_{12} \\ 0 & H_{12}^* & H_{22}(K) \end{pmatrix}$$

with the entries

$$\begin{aligned} H_{00}(K) f_0 &= w_0(K) f_0, \quad H_{01} f_1 = \int_{\mathbb{T}} v(t) f_1(t) dt, \\ (H_{11}(K) f_1)(x) &= w_1(K; x) f_1(x), \quad (H_{12} f_2)(x) = \int_{\mathbb{T}} v(t) f_2(x, t) dt, \\ (H_{22}(K) f_2)(x, y) &= w_2(K; x, y) f_2(x, y), \quad f_i \in H_i, \quad i = 0, 1, 2 \end{aligned}$$

where H_{ij}^* ($i < j$) denotes the adjoint operator to H_{ij} .

Here $w_0(\cdot)$ is a real-valued bounded function on $\mathbb{T}$, the function $v(\cdot)$ is a real-valued analytic on $\mathbb{T}$, the functions $w_1(\cdot; \cdot)$ and $w_2(\cdot; \cdot, \cdot)$ are defined by the equalities

$$w_1(K; x) := l_1 \varepsilon(x) + l_2 \varepsilon(K - x) + 1, \quad w_2(K; x, y) := l_1 \varepsilon(x) + l_1 \varepsilon(y) + l_2 \varepsilon(K - x - y),$$

respectively, with $l_1, l_2 > 0$ and

$$\varepsilon(x) := 1 - \cos(nx), \quad n \in \mathbb{N}.$$

Under these assumptions the operator $H(K)$ is bounded and self-adjoint.

We remark that the operators H_{01} and H_{12} , resp. H_{01}^* and H_{12}^* are called annihilation resp. creation operators, respectively. It is clear that

$$\begin{aligned} H_{01}^* : H_0 &\rightarrow H_1, \quad (H_{01}^* f_0)(x) = v(x) f_0, \quad f_0 \in H_0; \\ H_{12}^* : H_1 &\rightarrow H_2, \quad (H_{12}^* f_1)(x, y) = \frac{1}{2} (v(x) f_1(y) + v(y) f_1(x)), \quad f_1 \in H_1. \end{aligned}$$

To study the essential spectrum of the operator $H(K)$ we introduce a family of bounded self-adjoint operators (generalized Friedrichs models) $h(k)$, $k \in \mathbb{T}$, which acts in $H_0 \oplus H_1$ as

$$h(k) := \begin{pmatrix} h_{00}(k) & h_{01} \\ h_{01}^* & h_{11}(k) \end{pmatrix},$$

where

$$\begin{aligned} h_{00}(k) f_0 &= (l_2 \varepsilon(k) + 1) f_0, \quad h_{01} f_1 = \frac{1}{\sqrt{2}} \int_{\mathbb{T}} v(t) f_1(t) dt, \\ (h_{11}(k) f_1)(y) &= E_k(y) f_1(y), \quad E_k(y) := l_1 \varepsilon(y) + l_2 \varepsilon(k - y). \end{aligned}$$

Let the operator $h_0(k)$, $k \in \mathbb{T}$ acts in $H_0 \oplus H_1$ as

$$h_0(k) := \begin{pmatrix} 0 & 0 \\ 0 & h_{11}(k) \end{pmatrix}.$$

The perturbation $h(k) - h_0(k)$ of the operator $h_0(k)$ is a self-adjoint operator of rank 2, and thus, according to the Weyl theorem, the essential spectrum of the operator $h(k)$ coincides with the essential spectrum of $h_0(k)$. It is evident that $\sigma_{\text{ess}}(h_0(k)) = [E_{\min}(k); E_{\max}(k)]$, where the numbers $E_{\min}(k)$ and $E_{\max}(k)$ are defined by

$$E_{\min}(k) := \min_{y \in T} E_k(y) \quad \text{and} \quad E_{\max}(k) := \max_{y \in T} E_k(y).$$

This yields $\sigma_{\text{ess}}(h(k)) = [E_{\min}(k); E_{\max}(k)]$.

For any $k \in T$ we define an analytic function $\Delta(k; \cdot)$ (the Fredholm determinant associated with the operator $h(k)$ in $\square \setminus [E_{\min}(k); E_{\max}(k)]$) by

$$\Delta(k; z) := l_2 \varepsilon(k) + 1 - z - \frac{1}{2} \int_T \frac{v^2(t) dt}{E_k(t) - z}.$$

Note that for the discrete spectrum of $h(k)$ the equality

$$\sigma_{\text{disc}}(h(k)) = \{z \in \square \setminus [E_{\min}(k); E_{\max}(k)] : \Delta(k; z) = 0\}$$

holds (see Lemma 1).

The following theorem [5, 6] describes the location of the essential spectrum of the operator $H(K)$ by the spectrum of the family $h(k)$ of generalized Friedrichs models.

Theorem 1. For the essential spectrum of $H(K)$ the equality

$$\sigma_{\text{ess}}(H(K)) = \bigcup_{x \in T} \{\sigma_{\text{disc}}(h(K-x)) + l_1 \varepsilon(x)\} \cup [m_K; M_K] \quad (1)$$

holds, where the numbers m_K and M_K are defined by

$$m_K := \min_{x, y \in T} w_2(K; x, y) \quad \text{and} \quad M_K := \max_{x, y \in T} w_2(K; x, y).$$

Let Λ be a subset of T given by

$$\Lambda := \left\{0; \pm \frac{2}{n} \pi; \pm \frac{4}{n} \pi; \dots; \pm \frac{n'}{n} \pi\right\} \cup \Pi_n,$$

where

$$n' := \begin{cases} n-2, & \text{if } n \text{ is even} \\ n-1, & \text{if } n \text{ is odd} \end{cases} \quad \text{and} \quad \Pi_n := \begin{cases} \{\pi\}, & \text{if } n \text{ is even} \\ \emptyset, & \text{if } n \text{ is odd} \end{cases}$$

Direct calculation shows that the cardinality of Λ is equal to n . It is easy to check that for any $K \in \Lambda$ the function $w_2(K; \cdot, \cdot)$ has non-degenerate zero minimum at the points of $\Lambda \times \Lambda$, that is, $m_K = 0$ for $K \in \Lambda$.

Since $0 \in \Lambda$ the definition of the functions $w_1(\cdot; \cdot)$ and $w_2(\cdot; \cdot, \cdot)$ imply the identity $h(0) \equiv h(k)$ for all $k \in \Lambda$.

For $\delta > 0$ and $a \in T$ we set

$$U_\delta(a) := \{x \in T : |x - a| < \delta\}.$$

We remark that, if $v(x') = 0$ for some $x' \in T$, then from analyticity of $v(\cdot)$ on T it follows that there exist positive numbers C_1, C_2 and δ such that the inequalities

$$C_1 |x - x'|^{\beta(x')} \leq |v(x)| \leq C_2 |x - x'|^{\beta(x')}, \quad x \in U_\delta(x') \quad (2)$$

hold for some $\beta(x') \in N$.

Since for any $K \in \Lambda$ the function $w_2(K; \cdot, \cdot)$ has non-degenerate zero minimum at the points $(x', y') \in \Lambda \times \Lambda$, we obtain the following expansion

$$\begin{aligned} w_2(K; x, y) = & \frac{n^2}{2} \left[(l_1 + l_2) |x - x'|^2 + 2l_2(x - x', y - y') + (l_1 + l_2) |y - y'|^2 \right] \\ & + O(|x - x'|^4) + O(|y - y'|^4) \end{aligned}$$

as $|x - x'|, |y - y'| \rightarrow 0$ for $K, x', y' \in \Lambda$. Therefore, for any $K \in \Lambda$ there exist $C_1, C_2 > 0$ and $\delta > 0$ such that estimates hold

$$C_1(|x - x'|^2 + |y - y'|^2) \leq w_2(K; x, y) \leq C_2(|x - x'|^2 + |y - y'|^2), (x, y) \in U_\delta(x') \times U_\delta(y'); \quad (3)$$

$$w_2(K; x, y) \geq C_1, (x, y) \notin \bigcup_{x' \in \Lambda} U_\delta(x') \times \bigcup_{y' \in \Lambda} U_\delta(y'). \quad (4)$$

Hence, if $v(x') = 0$ for all $x' \in \Lambda$, then using the inequalities (2)–(4) one can easily see that the integral

$$\int_T \frac{v^2(t) dt}{w_2(K; x, t)}$$

is positive and finite for any $K \in \Lambda$ and $x \in T$.

For $x' \in \Lambda$ the Lebesgue dominated convergence theorem yields

$$\Delta(-x'; -l_1 \varepsilon(x')) = \lim_{x \rightarrow x'} \Delta(-x; -l_1 \varepsilon(x)),$$

and hence, if $v(x') = 0$ for all $x' \in \Lambda$, then the function $\Delta(-x; -l_1 \varepsilon(x))$ is a continuous with respect to x on T .

Set

$$\tau_{\text{ess}}(K) := \min\{\lambda : \lambda \in \sigma_{\text{ess}}(H(K))\}.$$

Then $\tau_{\text{ess}}(K) \in \sigma_{\text{ess}}(H(K))$ and it is called the lower bound of the essential spectrum of $H(K)$.

The main result of the present paper is the following theorem.

Theorem 2. Let $K \in \Lambda$. For the lower bound $\tau_{\text{ess}}(K)$ the following assertions hold.

- (i) If $v(x') \neq 0$ for some $x' \in \Lambda$, then $\tau_{\text{ess}}(K) < 0$;
- (ii) If $v(x') = 0$ for all $x' \in \Lambda$ and $\min_{x \in T} \Delta(-x; -l_1 \varepsilon(x)) < 0$ then $\tau_{\text{ess}}(K) < 0$;
- (iii) If $v(x') = 0$ for all $x' \in \Lambda$ and $\min_{x \in T} \Delta(-x; -l_1 \varepsilon(x)) \geq 0$, then $\tau_{\text{ess}}(K) = 0$.

This theorem is play a key role in the study of the number of eigenvalues of $H(K)$, $K \in \Lambda$.

Lower bound of the essential spectrum of $H(K)$

In this section first we study the discrete spectrum of the generalized Friedrichs model $h(k)$ and then we prove Theorem 2.

The following lemma is a simple consequence of the Birman-Schwinger principle and the Fredholm theorem.

Lemma 1. For any fixed $k \in T$ the operator $h(k)$ has an eigenvalue $z(k) \in \mathbb{R} \setminus [E_{\min}(k); E_{\max}(k)]$ if and only if $\Delta(k; z(k)) = 0$.

In the next two lemmas we describe the number and location of the eigenvalues of $h(k)$.

Lemma 2. If $v(y') \neq 0$ for some $y' \in \Lambda$, then for all $x' \in \Lambda$ the operator $h(x')$ has a unique negative eigenvalue.

Proof. Since the function $w_2(0; \cdot, \cdot)$ has the non-degenerate zero global minimum at the points $(0, 0) \in T^2$, by the implicit function theorem there exist $\delta > 0$ and an analytic function $y_0(\cdot)$ on $U_\delta(0)$ such that for any $x \in U_\delta(0)$ the point $y_0(x)$ is the unique non-degenerate minimum of the function $w_2(x, \cdot)$ and $y_0(0) = 0$. Therefore, we have $w_2(x, y_0(x)) = m_x$ for any $x \in U_\delta(0)$.

Let $w_2(\cdot, \cdot)$ be the function defined on $U_\delta(x) \times T$ as $w_2(x, y) := w_2(x, y + y_0(x))$. Then for any $x \in U_\delta(x)$ the function $w_2(x, \cdot)$ has a unique non-degenerate zero minimum at the point $x \in T$. Now using the equality

$$\int_T \frac{v_1^2(t) dt}{w_2(x, t) - m_x} = \int_T \frac{v_1^2(t + y_0(x)) dt}{w_2(x, t)}, \quad x \in U_\delta(0),$$

the continuity of the function $v_1(\cdot)$, the conditions $v_1(0) \neq 0$ and $y_0(0) = 0$ it is easy to see that

$$\lim_{z \rightarrow m_x - 0} \Delta(x; z) = -\infty \text{ for all } x \in U_\delta(0).$$

Since for any $x \in T$ the function $\Delta(x; \cdot)$ is continuous and monotonically decreasing on $(-\infty, m_x)$ the equality

$$\lim_{z \rightarrow -\infty} \Delta(x; z) = \infty \quad (5)$$

implies that for any $x \in U_\delta(0)$ the function $\Delta(x; \cdot)$ has a unique zero $z = z(x)$, lying in $(-\infty, m_x)$. By Lemma 1 the number $z(x)$ is the eigenvalue of $h(x)$.

Lemma 3. Let $v_1(x') = 0$, for any $x' \in \Lambda$.

(i) If $\min_{x \in T} \Delta(x; m) \geq 0$, then for any $x \in T$ the operator $h(x)$ has no eigenvalues, lying on the left of m .

(ii) If $\min_{x \in T} \Delta(x; m) < 0$, then there exists a non-empty set $G \subset T$ such that for any $x \in G$ the operator $h(x)$ has a unique eigenvalue $z(x)$, lying on the left of m .

Proof. First we recall that if $v_1(0) = 0$, then the function $\Delta(\cdot; m)$ is a continuous on T . Let $\min_{x \in T} \Delta(x; m) \geq 0$. Since for any $x \in T$ the function $\Delta(x; \cdot)$ is monotonically decreasing on $(-\infty, m)$ we have $\Delta(x; z) > \Delta(x; m) \geq \min_{x \in T} \Delta(x; m) \geq 0$, that is, $\Delta(x; z) > 0$ for all $x \in T$ and $z < m$. Therefore, by Lemma 1 for any $x \in T$ the operator $h(x)$ has no eigenvalues in $(-\infty, m)$.

Now we suppose that $\min_{x \in T} \Delta(x; m) < 0$ and introduce the following subset of T :

$$G := \{x \in T : \Delta(x; m) < 0\}.$$

Since $\Delta(\cdot; m)$ is a continuous on the compact set T , there exists a point $x^0 \in T$ such that $\min_{x \in T} \Delta(x; m) = \Delta(x^0; m)$, that is, $x^0 \in G$. So, the set G is a non-empty. Note that if $\max_{x \in T} \Delta(x; m) < 0$, then $\Delta(x; m) < 0$ for all $x \in T$ and hence $G = T$.

Since for any $x \in T$ the function $\Delta(x; \cdot)$ is a continuous and monotonically decreasing on $(-\infty, m]$ by the equality (5) for any $x \in G$ there exists a unique point $z(x) \in (-\infty, m)$ such that $\Delta(x; z(x)) = 0$. By Lemma 1 for any $x \in G$ the point $z(x)$ is the unique eigenvalue of $h(x)$.

By the construction of G the inequality $\Delta(x; m) \geq 0$ holds for all $x \in T \setminus G$. In this case one can see that for any $x \in T \setminus G$ the operator $h(x)$ has no eigenvalues in $(-\infty, m)$.

Proof of Theorem 2. Let $v_1(0) \neq 0$. Then by Lemma 2 there exists $\delta > 0$ such that for any $x \in U_\delta(0)$ the operator $h(x)$ has a unique eigenvalue $z(x)$, lying on the left of m_x . In particular, $z(0) < m_0$. Since $m = \min_{x \in T} m_x = m_0$ it follows that $\min \sigma \leq z(0) < m$, that is, $E_{\min} < m$.

Let $v_1(0) = 0$. Then two cases are possible: $\min_{x \in T} \Delta(x; m) \geq 0$ or $\min_{x \in T} \Delta(x; m) < 0$. In the case $\min_{x \in T} \Delta(x; m) \geq 0$, by the part (i) of Lemma 3 for any $x \in T$ the operator $h(x)$ has no eigenvalues in $(-\infty, m)$, that is, $\min \sigma \geq m$. By Theorem 1 it means that $E_{\min} = m$.

For the case $\min_{x \in T} \Delta(x; m) < 0$, using the part (ii) of Lemma 3 we obtain $\min \sigma \leq z(x') < m$ for all $x' \in G$, that is, $E_{\min} < m$.

REFERENCES:

1. C.Tretter. *Spectral Theory of Block Operator Matrices and Applications*. Imperial College Press, 2008.
2. A.Jeribi. *Spectral Theory of Block Operator Matrices and Applications*. Imperial College Press, 2008.
3. Muminov M., Neidhardt H., Rasulov T. *On the spectrum of the lattice spin-boson Hamiltonian for any coupling: 1D case*. *Journal of Mathematical Physics*, 2015, Vol. 56, 053507.
4. Rasulov T.Kh. *Branches of the essential spectrum of the lattice spin-boson model with at most two photons*. *Theoretical and Mathematical Physics*, 2016, Vol. 186, no. 2, pp. 251-267.
5. S.Albeverio, S.N.Lakaev, T.H.Rasulov. *On the spectrum of an Hamiltonian in Fock space. Discrete spectrum asymptotics*. *J. Stat. Phys.*, 127:2 (2007), pp. 191–220.
6. S.Albeverio, S.N.Lakaev, T.H.Rasulov. *The Efimov effect for a model operator associated with the Hamiltonian of a non conserved number of particles*. *Methods Funct. Anal. Topology*, 13:1 (2007), pp. 1–16.